

COLISTINE 210 MCI

Oral colistin solution for poultry and ruminants

Oral solution

COMPOSITION:

Colistin (as sulfate)210 MUI

Excipients s.q.f.100 ml

INDICATIONS:

In poultry and ruminants:

Treatment of infections induced by sensitive germs to colistin, mainly enteritis due to Salmonella and E.Coli.

DOSAGE AND ADMINISTRATION ROUTE:

Oral route, in drinking water.

- Poultry: 75 000 IU of colistin per Kg BW per day i.e. 0,25 ml of COLISTINE 210 MCI per L of drinking water for 3 consecutive days.

- Calves and lambs: 50 000 IU of colistin per Kg BW every 12 hours, i.e. 0.25 ml of COLISTINE 210 MCI per 10 Kg BW, twice a day, for 3 days. To mix with milk or with drinking water.

PRECAUTIONS FOR USE:

None.

CONTRA-INDICATIONS:

None.

SIDE EFFECTS:

None.

STORAGE:

Keep in its commercial packaging, at a temperature below 25°C.

WITHDRAWAL PERIOD:

Meat and offals: 7 days.

PACK SIZES:

Bottle of 250 ml

Bottle of 500 ml

Bottle of 1 L

Can of 2.5 L

Can of 5 L.

For Veterinary Use.

M.C.I.
Santé Animale



M.C.I. Lot 157, Z.I. Sud Ouest P.O. Box 278 - 28 810 MOHAMMEDIA - MOROCCO-
Phone: + 212 5 23 30 31 32+ Fax : + 212 5 23 30 21 30- Web Site: www.mci-santeanimale.com