

GANIDAN

**Anti-diarrheal for cattle, sheep, goats, equines, dogs, cats,
poultry and rabbits**

Oral powder

COMPOSITION:

Sulfaguanidine9 g
Excipient s.q.f.10 g

INDICATIONS:

Treatment of the infectious enteritis and gastroenteritis caused by germs susceptible to sulfaguanidine.

Treatment of the coccidiosis of poultry and rabbits.

DOSAGE AND ADMINISTRATION ROUTE:

Oral route, according to the weight:

Adult cattle and equine.....30 g to 40 g
Calves and foals8 g to 10 g
Goats and sheep.....4 g to 8 g
Pics.....4 g to 10 g

Administer once or twice a day and reduce the posology from the 3rd day. Treat for 4 to 8 days.

Poultry and rabbits:

GANIDAN should be administered mixed with dry ration in the proportions of 0.5 to 1%, during 3 to 5 days.

The animals should be well watered during the treatment period.

PRECAUTION OF USE:

In animals: it's necessary to verify the absence of ulcerative colitis and dehydration of the animal before administration.

In the perso giving the product it's recommended to a mask, glasses and gloves.

WITHDRAWAL PERIOD:

Meat : 7 days,
Milk : 4 days,
Eggs : do not use in animals which eggs are intended for human consumption.

STORAGE:

Keep in its commercial packaging, in a cool and dry place.

M.C.I.
Santé Animale



M.C.I. Lot 157, Z.I. Sud Ouest P.O. Box 278 - 28 810 MOHAMMEDIA - MOROCCO-
Phone: + 212 5 23 30 31 32+ Fax : + 212 5 23 30 21 30- Web Site: www.mci-santeanimale.com