

# MULTICYNE

## Oral enrofloxacin and colistin for poultry

Oral solution

### COMPOSITION:

Colistin sulfate .....	200 MUI
Enrofloxacin .....	10 g
Excipients s.q.f. ....	100 ml

### INDICATIONS:

In poultry:

- Prevention and treatment of Chronic Respiratory Disease (CRD) due to *Haemophilus*, *Pasteurella* and *Mycoplasma*.
- Prevention and treatment of infectious gastro-enteritis as *Collibacillosis* and *Salmonellosis*.

### DOSAGE AND ADMINISTRATION ROUTE:

Oral route, in drinking water.

Colistin: 50 000 to 100 000 IU per Kg of BW per day, Enrofloxacin: 5 to 10 mg per Kg of B.W per day, corresponding to 0.5 ml of MULTICYNE per L of drinking water during 3 to 5 days. Adjust the posology to the water consumption.

### PRECAUTIONS FOR USE:

- Medicated water must be consumed within 24 hours after reconstitution
- Respect the prescribed doses and the treatment duration.

### CONTRA-INDICATIONS:

Do not administer to layers when eggs are intended to human consumption.

### SIDE EFFECTS:

None.

### STORAGE:

Keep in its commercial packaging, at a temperature below 25°C.

### WITHDRAWAL PERIOD:

Meat and offals: 7 days.

### PACK SIZES:

- Bottle of 250 ml
- Bottle of 500 ml
- Bottle of 1 L
- Can of 2.5 L
- Can of 5 L.

For Veterinary Use.

**M.C.I.**  
**Santé Animale**



M.C.I. Lot 157, Z.I. Sud Ouest P.O. Box 278 - 28 810 MOHAMMEDIA - MOROCCO-  
Phone: + 212 5 23 30 31 32+ Fax : + 212 5 23 30 21 30- Web Site: [www.mci-santeanimale.com](http://www.mci-santeanimale.com)