

# TYLOSINE MCI

Oral antibiotic for poultry

Oral powder

## COMPOSITION:

Tylosine ( as tartrate)..... 100 %

## INDICATIONS:

Treatment and prevention of avian mycoplasmosis:

- Mycoplasmosis primitive.
- Chronic Respiratory Diseases.

Prevention and treatment of infectious sinusitis in turkeys.

Control of calf pneumonia caused by *Pasteurella multocida* and Mycoplasma.

## DOSAGE AND ADMINISTRATION ROUTE :

*Chickens and Turkeys:*

- Curative treatment: 0.5 g / liter of drinking water for 3-5.
- Preventive treatment: 0.5 g / liter of drinking water for the first three days of life and during 24 hours at the 4th week or during stress (vaccinations, etc.).

To obtain medicated water containing 0.5 g of tylosin per liter, 100 g of TYLOSINE MCI should be diluted in 250 mL of warm water then shake well for complete dissolution of the preparation. Mix this solution with 200 liters of water.

*Calves:* Administer 1 g of Tylosin per animal 2 times a day for 2 weeks at the entry in nursery.

## CONTRA-INDICATIONS:

Do not administer to rabbits.

## STORAGE:

Keep in a dry place in a temperature below 25° C.

## WITHDRAWAL TIME:

Chicken: Meat and offal: 2 days / eggs: 5 days.

Turkey: Meat and offal: 3 days.

Calf: Meat and offal: 10 days.

## PACK SIZES:

Pot of 100 g.

**M.C.I.**  
**Santé Animale**



M.C.I. Lot 157, Z.I. Sud Ouest P.O. Box 278 - 28 810 MOHAMMEDIA - MOROCCO-  
Phone: + 212 5 23 30 31 32+ Fax: + 212 5 23 30 21 30- Web Site: [www.mci-santeanimale.com](http://www.mci-santeanimale.com)