

BENDAR BOLUS 2500

Internal anthelmintic for cattle, dromedaries and equines

Bolus

COMPOSITION:

Each 15 000 mg bolus contains:
Albendazole 2500 mg

INDICATIONS:

In cattle, dromedaries and equines:
Treatment and prophylaxis of worm-infections caused by nematodes, cestodes and trematodes of the gastrointestinal tract, respiratory tract and liver, at both adult and larval states.

DOSAGE AND ADMINISTRATION ROUTE:

Oral route.

Cattle:

- 1 bolus / 350 Kg BW, corresponding approximately to 7 mg of albendazole/Kg BW for gastro-intestinal and lung strongyles and taeniasis, corresponding to 1 bolus for a cattle of 350 Kg BW.
- 1 bolus / 250 Kg BW, corresponding approximately to 10 mg of albendazole/Kg BW for trematodes, corresponding to 1,5 bolus for a cattle of 400 Kg BW.

Equines and dromedaries:

- 1 bolus / 360 Kg BW, corresponding approximately to 7 mg of albendazole/Kg BW for gastrointestinal and lung strongyles and taeniasis, corresponding to 2 bolus for a camel around 800 Kg BW and 1.5 bolus for an equine of 600 Kg BW.

PRECAUTIONS FOR USE:

None.

CONTRA-INDICATIONS:

- Do not use for pregnant cows during the first month of pregnancy as a precaution of a possible risk on embryonic development.
- Do not use in dairy females when milk is intended for human consumption.

SIDE EFFECTS:

None.

STORAGE:

Keep in its commercial packaging, at a temperature below 25°C.

WITHDRAWAL PERIOD:

Meat and offals: 14 days.

Milk: See the contra-indications.

PACK SIZES:

- Box of 24 Bolus
- Box of 50 Bolus
- Box of 100 Bolus
- Box of 200 Bolus.

For Veterinary Use.

M.C.I.
Santé Animale



M.C.I. Lot 157, Z.I. Sud Ouest P.O. Box 278 - 28 810 MOHAMMEDIA - MOROCCO-
Phone: + 212 5 23 30 31 32+ Fax : + 212 5 23 30 21 30- Web Site: www.mci-santeanimale.com